

CHEMICAL BONDING

Chapter

5

IMRAN SYAKIR BIN MOHAMAD | MOHD HAIZAL BIN MOHD HUSIN
FACULTY OF MECHANICAL TECHNOLOGY AND ENGINEERING

Valence electrons are the outer shell electrons of an atom. The valence electrons are the electrons that participate in chemical bonding.

<u>Group</u>	<u>e⁻ configuration</u>	<u># of valence e⁻</u>
1A	ns ¹	1
2A	ns ²	2
3A	ns ² np ¹	3
4A	ns ² np ²	4
5A	ns ² np ³	5
6A	ns ² np ⁴	6
7A	ns ² np ⁵	7



Lewis Dot Symbols

1 1A	2 2A											13 3A	14 4A	15 5A	16 6A	17 7A	18 8A
·H·												·B·	·C·	·N·	·O·	·F·	He:
·Li·	·Be·											·Al·	·Si·	·P·	·S·	·Cl·	·Ne·
3 3B	4 4B	5 5B	6 6B	7 7B	8 8B	9 8B	10 8B	11 1B	12 2B								
·K·	·Ca·									·Ga·	·Ge·	·As·	·Se·	·Br·	·Kr·		
·Rb·	·Sr·									·In·	·Sn·	·Sb·	·Te·	·I·	·Xe·		
·Cs·	·Ba·									·Tl·	·Pb·	·Bi·	·Po·	·At·	·Rn·		
·Fr·	·Ra·																

The Ionic Bond (Electrovalence)


An ionic bond is the electrostatic force that holds ions together in an ionic compound

Ionic compound combine a Group IA & Group IIA metal with a halogen or oxygen




The Covalent Bond


A **covalent bond** is a chemical bond in which two or more electrons are shared by two atoms. (Non metal & non metal)




Why should two atoms share electrons?




Lewis structure of F_2




Lewis structure of water





Double bond – two atoms share two pairs of electrons



Triple bond – two atoms share three pairs of electrons



Lengths of Covalent Bonds



Bond Lengths

Triple bond < Double Bond < Single Bond

Bond Type	Bond Length (pm)
C-C	154
C=C	133
C \equiv C	120
C-N	143
C=N	138
C \equiv N	116



Comparison of Ionic and Covalent Compounds

Table 9.3 Comparison of Some General Properties of an Ionic Compound and a Covalent Compound

Property	NaCl	CCl ₄
Appearance	White solid	Colorless liquid
Melting point (°C)	801	−23
Molar heat of fusion* (kJ/mol)	30.2	2.5
Boiling point (°C)	1413	76.5
Molar heat of vaporization* (kJ/mol)	600	30
Density (g/cm ³)	2.17	1.59
Solubility in water	High	Very low
Electrical conductivity		
Solid	Poor	Poor
Liquid	Good	Poor

* Molar heat of fusion and molar heat of vaporization are the amounts of heat needed to melt 1 mole of the solid and to vaporize 1 mole of the liquid, respectively.


Polar covalent bond or ***polar bond*** is a covalent bond with greater electron density around one of the two atoms



Electronegativity is the ability of an atom to attract toward itself the electrons in a chemical bond.

Electron Affinity - **measurable**, Cl is highest

Electronegativity - **relative**, F is highest



Both are related but different concepts. EA refers to an isolated atom and E refers to an atom in chemical bond. Usually, $EA > E$.

Electronegativities of Common Elements

Increasing electronegativity

Increasing electronegativity

1A																	8A	
H 2.1	2A												3A	4A	5A	6A	7A	
Li 1.0	Be 1.5												B 2.0	C 2.5	N 3.0	O 3.5	F 4.0	
Na 0.9	Mg 1.2												Al 1.5	Si 1.8	P 2.1	S 2.5	Cl 3.0	
K 0.8	Ca 1.0	3B	4B	5B	6B	7B	8B			1B	2B		Ga 1.6	Ge 1.8	As 2.0	Se 2.4	Br 2.8	
Rb 0.8	Sr 1.0	Sc 1.3	Ti 1.5	V 1.6	Cr 1.6	Mn 1.5	Fe 1.8	Co 1.9	Ni 1.9	Cu 1.9	Zn 1.6		In 1.7	Sn 1.8	Sb 1.9	Te 2.1	I 2.5	
Cs 0.7	Ba 0.9	La-Lu 1.0-1.2	Hf 1.3	Ta 1.5	W 1.7	Re 1.9	Os 2.2	Ir 2.2	Pt 2.2	Au 2.4	Hg 1.9		Tl 1.8	Pb 1.9	Bi 1.9	Po 2.0	At 2.2	
Fr 0.7	Ra 0.9																	

Classification of bonds by difference in electronegativity

Difference

Bond Type

0

Covalent

≥ 2

Ionic

$0 < \text{and} < 2$

Polar Covalent

Increasing difference in electronegativity



Covalent

Polar Covalent

Ionic



share e^-

partial transfer of e^-

transfer e^-



Classify the following bonds as ionic, polar covalent, or covalent: The bond in CsCl ; the bond in H_2S ; and the NN bond in H_2NNH_2 .

Intermolecular Forces

Intermolecular forces are attractive forces **between** molecules.

Intramolecular forces hold atoms together in a molecule.

Intermolecular vs Intramolecular

- 41 kJ to vaporize 1 mole of water (**inter**)
- 930 kJ to break all O-H bonds in 1 mole of water (**intra**)



Generally,
intermolecular
forces are much
weaker than
intramolecular
forces.

“Measure” of intermolecular force

boiling point

melting point


ΔH_{vap}

Intermolecular Forces

Dipole-Dipole Forces

Attractive forces between **polar molecules**

Orientation of Polar Molecules in a Solid




Intermolecular Forces

Ion-Dipole Forces

Attractive forces between an **ion** and a **polar molecule**


Ion-Dipole Interaction




Intermolecular Forces

Dispersion Forces (London)

Attractive forces that arise as a result of **temporary dipoles induced** in atoms or molecules



ion-induced dipole interaction



dipole-induced dipole interaction

Intermolecular Forces

Dispersion Forces Continued

Polarizability is the ease with which the electron distribution in the atom or molecule can be distorted.

Polarizability increases with:

- greater number of electrons
- more diffuse electron cloud



Dispersion forces usually increase with molar mass.

Table 11.2 Melting Points of Similar Nonpolar Compounds

Compound	Melting Point (°C)
CH ₄	-182.5
CF ₄	-150.0
CCl ₄	- 23.0
CBr ₄	90.0
Cl ₄	171.0

What type(s) of intermolecular forces exist between each of the following molecules?

HBr

CH₄

SO₂


Intermolecular Forces

Hydrogen Bond

The **hydrogen bond** is a special dipole-dipole interaction between the hydrogen atom in a polar N-H, O-H, or F-H bond and an electronegative O, N, or F atom.



A & B are N, O, or F



Why is the hydrogen bond considered a “special” dipole-dipole interaction?

